

Opportunistic diseases among HIV-infected patients: a multicenter-nationwide Korean HIV/AIDS cohort study, 2006 to 2013

Youn Jeong Kim¹, Jun Hee Woo², Min Ja Kim³, Dae Won Park³, Joon-Young Song³, Shin Woo Kim⁴, Jun Yong Choi⁵, June Myung Kim⁵, Sang Hoon Han⁵, Jin-Soo Lee⁶, Bo Youl Choi⁷, Joo Shil Lee⁸, Sung-Soon Kim⁸, Mee-Kyung Kee⁸, Moon Won Kang¹, and Sang Il Kim¹

¹Division of Infectious Disease, Department of Internal Medicine, College of Medicine, Seoul St. Mary's Hospital, The Catholic University of Korea, Seoul; ²Department of Infectious Disease, Asan Medical Center, University of Ulsan College of Medicine, Seoul; ³Department of Internal Medicine, Korea University College of Medicine, Seoul; ⁴Department of Internal Medicine, Kyungpook National University School of Medicine, Daegu; ⁵Department of Internal Medicine, Yonsei University College of Medicine, Seoul; ⁶Department of Internal Medicine, Inha University School of Medicine, Incheon; ⁷Department of Preventive Medicine, Hanyang University College of Medicine, Seoul; ⁸Korea Centers for Disease Control and Prevention, Cheongju, Korea

Received: October 22, 2014

Revised: January 6, 2015

Accepted: April 27, 2015

Correspondence to
Sang Il Kim, M.D.

Division of Infectious Disease,
Department of Internal
Medicine, College of Medicine,
Seoul St. Mary's Hospital, The
Catholic University of Korea, 222
Banpo-daero, Seocho-gu, Seoul
06591, Korea
Tel: +82-2-2258-1240
Fax: +82-2-2258-1254
E-mail: drksi@catholic.ac.kr

Background/Aims: The frequencies of opportunistic diseases (ODs) vary across countries based on genetic, environmental, and social differences. The Korean HIV/AIDS cohort study was initiated in 2006 to promote research on human immunodeficiency virus (HIV) infection in Korea, and to provide a logistical network to support multicenter projects on epidemiological, clinical, and laboratory aspects of HIV infection. This study evaluated the prevalence of ODs among HIV-infected patients in the era of highly active antiretroviral therapy, and the risk factors associated with ODs.

Methods: The study enrolled 1,086 HIV-infected patients from 19 hospitals. This study examined the baseline data of the HIV/AIDS Korean cohort study at the time of enrollment from December 2006 to July 2013.

Results: Candidiasis was the most prevalent opportunistic infection (n = 176, 16.2%), followed by *Mycobacterium tuberculosis* infection (n = 120, 10.9%), *Pneumocystis jirovecii* pneumonia (n = 121, 11.0%), cytomegalovirus infection (n = 52, 4.7%), and herpes zoster (n = 44, 4.0%). The prevalence rates of Kaposi's sarcoma (n = 8, 0.7%) and toxoplasmosis (n = 4, 0.4%) were very low compared with other countries. The risk factors for ODs were a low CD4 T cell count at the time of HIV diagnosis (odds ratio [OR], 1.01; $p < 0.01$), current smoking (OR, 2.27; $p = 0.01$), current alcohol use (OR, 2.57; $p = 0.04$), and a history of tuberculosis (OR, 5.23; $p < 0.01$).

Conclusions: Using recent Korean nationwide data, this study demonstrated that an important predictor of ODs was a low CD4 T cell count at the time of HIV diagnosis. Tuberculosis remains one of the most important ODs in HIV-infected patients in Korea.

Keywords: AIDS-related opportunistic infections; HIV; Korea; Tuberculosis

INTRODUCTION

Globally, an estimated 34 million people were infected with human immunodeficiency (HIV) at the end of 2011. In Korea, after the first case of HIV infection was reported in 1985, the number of cumulative HIV survivors rose to approximately 7,000 at the end of 2011, although the overall prevalence in the population was very low (< 1%) [1]. HIV infection leads to immunosuppression, which can result in life-threatening opportunistic diseases (ODs). Combination antiretroviral therapy reduced the mortality and morbidity [2,3], but despite advances in antiretroviral therapy, ODs remain one of the main causes of morbidity and mortality in HIV/AIDS patients.

The frequencies of ODs may vary across countries because of differences in genetic, environmental, and social backgrounds. Two studies have examined the epidemiology and clinical manifestations of ODs in Korean patients with HIV/AIDS [4,5]. The introduction of combination antiretroviral therapy and the use of prophylactic antimicrobials for ODs have changed the prevalence of ODs over time. Consequently, an understanding of the current epidemiology of ODs among HIV/AIDS patients should help to establish guidelines for the prevention of ODs. This study evaluated the prevalence of ODs among HIV-infected patients in the era of highly active antiretroviral therapy, and the risk factors associated with ODs.

METHODS

Study design and time period

The cohort recruited patients older than 18 years infected with HIV after obtaining informed consent. Trained researchers from all centers prospectively collected information every 6 months, using a standardized protocol. The information included the medical history, socioeconomic status, physical findings, laboratory findings including immunological and virological status, and ODs. In this study, we determined the prevalence of ODs using the baseline data of the HIV/AIDS cohort study at the time of enrollment, from December 2006 to July 2013.

Definition

ODs were diagnosed from the clinical, laboratory, and pathology findings, following published criteria [6]. A diagnosis of tuberculosis (TB) was based on compatible symptoms/signs and positive acid-fast bacilli (AFB) smear, culture, polymerase chain reaction (PCR), and histological or radiological findings compatible with TB. Disseminated TB was defined when more than two sites were involved [7]. *Pneumocystis jirovecii* pneumonia was confirmed based on compatible symptoms/signs, and radiological findings such as bilateral, diffuse interstitial infiltrates, demonstration of the cyst wall on methenamine silver staining, or a positive *P. jirovecii* PCR. Cytomegalovirus (CMV) viremia was detected using a CMV PCR assay. CMV retinitis was diagnosed by an ophthalmologist. CMV disease such as pneumonia or hepatitis was diagnosed based on compatible symptoms and pathology staining for CMV.

Ethics approval

Ethics approval was obtained from all participating hospitals.

Data analysis

Data are provided as mean \pm standard deviation, or as proportions. Student *t* test or the Mann-Whitney *U* test was used to analyze continuous variables, and the chi-square test or Fisher exact test was used for categorical variables. We analyzed the risk factors associated with developing ODs in HIV-infected patients using both univariate and multivariate logistic regression analyses. Variables with a *p* value < 0.1 in the univariate analysis were added in a forward stepwise manner and selected to create the final model for multivariate analysis with forward variables. Data were analyzed using the SPSS version 16.0 (SPSS Inc., Chicago, IL, USA), and a *p* < 0.05 was considered statistically significant.

RESULTS

The study enrolled 1,086 HIV-infected patients (1,007 males and 79 females; median age, 41 years) at 19 university hospitals from 2006 to 2012 (Table 1). The majority of patients (*n* = 601, 55.4%) were 30 to 50 years of age. The median CD4⁺ T cell count at enrollment was 352 cells/

Table 1. Comparisons of demographic characteristics between patients with ODs and without ODs

Characteristic	Total (n = 1,086)	ODs (+) (n = 297)	ODs (-) (n = 789)	p value
Age, yr	41 (33–50)	44 (36–52)	41 (32–50)	< 0.01
Male sex	1,007 (92.7)	279 (93.9)	728 (92.3)	0.37
Transmission route				
Sexual contact	945 (97.0) ^a	261 (87.8) ^a	684 (86.7) ^a	0.89
Transfusion or blood product	24 (2.5) ^a	9 (3.0) ^a	15 (1.9) ^a	0.87
Injection drug use	5 (0.5) ^a	3 (1.0) ^a	2 (0.3) ^a	0.09
Vertical transmission	0	0	0	-
CD4 T cell count at enrollment, cell/mm ³	352 (207–514)	240 (111–395)	388 (254–560)	< 0.01
CD4 T cell at HIV diagnosis, cell/mm ³	237 (93–377)	75 (25–203)	279.5 (169–420)	< 0.01
Viral load at enrollment, log copies/mL	5.69 (3.88–6.24)	5.04 (4.17–5.56)	4.57 (3.8–6.09)	0.62
HARRT at enrollment	730 (67.3)	232 (78.1)	498 (63.1)	< 0.01
Body mass index, kg/m ²	21.5 (19.4–23.5)	21.1 (19–23.5)	21.6 (19.8–23.6)	0.77
Smoking				
Current	465 (71.5) ^a	111 (59.4)	354 (76.5)	< 0.01
Former	648 (68.0) ^a	189 (70.0)	459 (67.2)	0.40
Alcohol				
Current	483 (73.0) ^a	106 (57.9)	377 (78.7)	< 0.01
Former	656 (70.0) ^a	183 (68.3)	473 (70.0)	0.47
Tuberculosis history	196 (18.0)	115 (38.7)	81 (10.2)	< 0.01
Hypertension	112 (10.3)	29 (9.7)	83 (10.5)	0.84
Diabetes mellitus	71 (6.5)	17 (5.7)	54 (6.8)	0.38
Previous BCG history	377 (54.5) ^a	108 (36.4)	269 (34.1)	0.71

Values are presented as median (interquartile range) or number (%).

OD, opportunistic disease; HIV, human immunodeficiency; HARRT, highly active antiretroviral therapy; BCG, Bacillus Calmette-Guérin.

^aThis data included only on that complete a questionnaire.

mm³ (interquartile range [IQR], 207 to 514), and 22.9% of the patients (n = 206) had fewer than 200 cells/mm³. The median viral load at enrollment was 5.69 log copies/mL (IQR, 3.88 to 6.24), while 53.2% of the patients had undetectable viral loads. At enrollment, 76% of the patients (n = 355) were receiving antiretroviral agents (Table 1).

Opportunistic diseases

Among the 1,086 HIV-infected patients, 613 ODs developed in 297 patients (24.4%), a rate of 2.1 episodes/patient. Table 2 summarizes the ODs seen in our study. Candidiasis was the most prevalent OD (n = 176, 16.2%), and was mainly oral candidiasis (n = 120), followed by esophageal candidiasis (n = 54). TB was the second most frequent OD (n = 120, 11.0%). Among the patients with

TB, 28.3% (n = 34) were taking antiretroviral treatment and 54.2% (n = 65), 15.8% (n = 19), and 15% (n = 18) had pulmonary, lymph node, and disseminated TB, respectively. The main diagnostic method for TB was the AFB smear (n = 58, 48.3%), followed by AFB culture (n = 49, 49%), PCR (n = 28, 23.3%), and pathology (n = 20, 16.6%). Based on clinical and radiological characteristics, anti-TB medication was given empirically to three patients with negative results on a routine test for TB. Of the 19 patients who had antimicrobial drug resistance tests, 26.3% (n = 5) were resistant to isoniazid, 15.8% (n = 3) to rifampin, and 10.5% (n = 2) to ethambutol. *P. jirovecii* pneumonia (n = 121, 11.1%) was the third most common OD, followed by CMV infection (n = 52, 4.7%), and herpes zoster (n = 44, 4.0%). CMV retinitis was found in 19

Table 2. Distribution of opportunistic disease and CD4 T cell count

Variable	No. of cases	CD4 T cell, cell/mm ³
Candidiasis	176 (16.2)	
Oral candidiasis	120	49 (17–133)
Esophageal candidiasis	54	43 (19–95.5)
Respiratory candidiasis	2	-
Tuberculosis	120 (11.0)	87 (37–205)
Nontuberculous Mycobacteria infection	2 (0.2)	-
<i>Pneumocystis jirovecii</i> pneumonia	121 (11.1)	30 (11–68)
CMV infection	52 (4.7)	25.5 (12.75–101.5)
Retinitis	19	
Pneumonia	13	
Hepatitis	9	
Colitis	2	
Cytomegalovirus viremia	9	
Herpes zoster	44 (4.0)	185 (116.5–273.5)
Herpes simplex virus infection	8 (0.7)	128 (10–320)
Recurrent pneumonia	7 (0.7)	62.5 (12–170)
Progressive multifocal leukoencephalopathy	6 (0.6)	107 (64.25–154)
Cryptococcosis	6 (0.6)	31.2 (9.5–76.75)
Toxoplasmosis	4 (0.4)	107 (25.5–246.25)
Isosporiasis	1 (0.09)	-
Non-Hodgkin's lymphoma	4 (0.4)	128 (26–222.5)
Burkitt lymphoma	1	
Diffuse large B-cell lymphoma	3	
Kaposi's sarcoma	8 (0.7)	133 (28–269)
Cervical carcinoma	1 (0.09)	-

Values are presented as number (%) or median (interquartile range).

patients. The prevalence rates of Kaposi's sarcoma (n = 8, 0.7%) and toxoplasmosis (n = 4, 0.4%) were low.

The median CD4 T cell count was 49 cells/mm³ (IQR, 17 to 133) in patients with esophageal candidiasis, 30 cells/mm³ (IQR, 11 to 68) in *P. jirovecii* pneumonia cases, and 87 cells/mm³ (IQR, 37 to 205) in TB cases (Table 2). When patients were first diagnosed with HIV, 13.9% (n = 151) had AIDS-defining illnesses. The most prevalent indicator of AIDS was *P. jirovecii* pneumonia (n = 91, 60.3%), followed by *Mycobacterium tuberculosis* infection (n = 69, 45.6%) and esophageal candidiasis (n = 38, 25.2%).

Fig. 1 shows the distribution of the CD4 cell count in study patients with OD. Patients who had a CD4 T cell count less than 50 cells/mm³ had a high prevalence of *P. jirovecii* pneumonia (66.9%), CMV infection (56.2%),

or esophageal or respiratory candidiasis (56.3%). A high prevalence of TB (73.8%) was found in patients with a CD4 T cell count less than 200 cells/mm³.

Risk factors for opportunistic diseases

Table 1 compares the demographic characteristics of patients with and without ODs. There were no differences in sex, transmission route, viral load, or body mass index. Patients with ODs were significantly older than patients without ODs (median, 44; IQR, 36 to 52 years vs. median, 41; IQR, 32 to 50 years; *p* < 0.01), and had a lower CD4 T cell count at the time of HIV diagnosis (median, 75; IQR, 25 to 203 cells/mm³ vs. median, 279.5; IQR, 169 to 420 cells/mm³, *p* < 0.01), and also had a lower CD4 T cell count at enrollment (median, 240; IQR, 111 to 395 cells/




Figure 1. Distribution of CD4 cell count in study patients with opportunistic disease. PCP, *Pneumocystis carinii* pneumonia; CMV, cytomegalovirus.

mm³ vs. median, 388; IQR, 254 to 560 cells/mm³, $p < 0.01$). The patients with ODs were more likely to be current smokers ($p < 0.01$), consume alcohol ($p < 0.01$), and have prior TB ($p < 0.01$) than those without ODs.

In the univariate analysis, older age (OR, 0.98; $p < 0.01$), a lower CD4 T cell count at the time of HIV diagnosis (OR, 1.01; $p < 0.01$), a lower CD4 T cell count at cohort study enrollment (OR, 1.01; $p < 0.01$), antiretroviral therapy (OR, 2.47; $p < 0.01$), current smoking (OR, 2.22; $p = 0.001$), current alcohol consumption (OR, 2.68; $p < 0.01$),

and a history of TB treatment (OR, 5.61; $p < 0.01$) were associated with ODs among HIV-infected patients (Table 3). The multivariate analysis showed that a low CD4 T cell count at the time of HIV diagnosis (OR, 1.01; $p < 0.01$), current smoking (OR, 2.27; $p = 0.01$), current alcohol consumption (OR, 2.57; $p = 0.04$), and a history of TB (OR, 5.23; $p < 0.01$) were independent risk factors for ODs.

Outcomes

During the follow-up period, 26 patients (2.4%) died. The ODs-related mortality was 0.7% ($n = 8$), accounting for 33.3% of deaths. The causes of OD-related death were pneumonia of unknown origin ($n = 4$), TB ($n = 1$), *P. jirovecii* pneumonia ($n = 1$), and *P. jirovecii* pneumonia combined with TB ($n = 1$).

DISCUSSION

Before the introduction of combination antiretroviral therapy, ODs were the main cause of death in HIV-infected patients. However, advances in antiviral treatment led to a dramatic decline in ODs [2,8,9]. In our study, although 24.4% of the HIV-infected patients developed ODs, they were not the main cause of death, unlike previous studies.

In our series, the most prevalent OD was candidiasis, followed by TB and *P. jirovecii* pneumonia. In a study of 173 HIV-infected patients performed from 1985 to 1998, TB was the most frequent opportunistic infection (25% of patients), followed by candidiasis and herpes zoster

Table 3. Risk factors for opportunistic disease among HIV patients

Factor	Univariate analysis			Multivariate analysis		
	OR	95% CI	p value	OR	95% CI	p value
Age	0.98	0.96–0.99	< 0.01	0.98	0.95–1.01	0.29
Body mass index	1.01	0.98–1.02	0.77			
CD4 T cell, HIV diagnosis	1.01	1.01–1.05	< 0.01	1.01	1.03–1.07	< 0.01
Viral load at enrollment	1	0.90–1.02	0.67			
CD4 T cell at enrollment	1.01	1.02–1.04	< 0.01	1.001	1.01–1.03	0.45
HAART at enrollment	2.47	1.79–3.41	< 0.01	1.41	0.55–3.56	0.47
Current smoking	2.22	1.55–3.19	< 0.01	2.27	1.22–4.26	0.01
Current alcohol	2.68	1.86–3.87	< 0.01	2.57	1.34–4.91	0.04
Tuberculosis history	5.61	4.03–7.7	< 0.01	5.23	2.69–10.2	< 0.01

HIV, human immunodeficiency; OR, odds ratio; CI, confidence interval; HAART, highly active antiretroviral therapy.

[5]. In another study of 176 HIV-infected patients from 1985 to 2000, candidiasis was the most prevalent OD, followed by *P. jirovecii* pneumonia and TB [4]. Based on our data, TB can be considered a re-emerging OD in the HIV population in Korea. The prevalence of ODs in HIV patients varies by location and is 32.5% in China and 27.4% in Ethiopia and New York city [10-12]. In developing countries, TB remains the most common cause of death among HIV-infected patients, similar to the situation in Korea, and unlike countries with low risks of TB [13,14]. In Korea, the estimated TB incidence in 2012 was 108 per 100,000 population, the highest among Organization for Economic Cooperation and Development countries. The identification and treatment of latent TB infection has lowered the incidence of TB in developed countries. The current guidelines for the management of TB in Korea recommend treating latent TB in HIV-infected patients [7]. Until now, many doctors in Korea have tended to avoid giving TB treatment for latent TB for several reasons: toxicity, drug interactions, and concern about drug resistance. In addition, our data showed that TB developed in patients with a low CD4 T cell count, and many studies demonstrated that impaired cellular immunity was associated with TB reactivation [15,16]. After the guidelines for latent TB are applied to clinical practice in Korea, further studies must examine the effect on the incidence of TB. In our series, a history of TB was associated with ODs among HIV-infected patients. Several studies have also demonstrated that a history of TB was usually associated with TB infection [17,18]. In intermediate TB endemic areas, TB reactivation is more common in HIV-infected patients than *de novo* TB. There were two cases of nontuberculous *Mycobacteria* infection in our study. Our study used the patient data at the time of enrollment, and further studies will examine prospectively collected cohort data.

Our study found a low prevalence of Kaposi's sarcoma and toxoplasmosis compared with other countries. This is because of the geographic variation in Kaposi's sarcoma. de Sanjose et al. [19] reported that the prevalence of Kaposi's sarcoma-associated herpesvirus in the general female population in Korea was only 4.93%.

The associated risk factors for ODs in our study were a low CD4 T cell count at the time of HIV diagnosis, smoking, alcohol consumption, and a history of TB. A low CD4 T cell count is associated with the develop-

ment of ODs [10,20]. A study of homeless HIV/AIDS patients in Korea showed that they are at an advanced stage at admission, which leads to ODs and an unfavorable outcome [21]. Our results showed that 97% of HIV-infected patients with *P. jirovecii* pneumonia had CD4 T cell counts less than 200 cells/mm³; this supports recent guidelines that patients with such low CD4+ T cell counts should receive prophylaxis for *P. jirovecii*. Smoking could influence inflammatory and immune responses in the oral cavity, and smoking was associated with oral candidiasis in some studies [22,23].

This study had several limitations. First, we included only patient data at study enrollment; prospective longitudinal data will provide more accurate data on OD incidence and change in ODs in Korea. Second, we did not calculate the risk factors for ODs for each disease, and did not describe the clinical course of each OD. Third, we included only the data from patients in our cohort study. However, this study enrolled patients from multiple centers that covered most areas of Korea, so it is likely to be representative epidemiological data for Korea.

In conclusion, there was a range of ODs among HIV-infected patients in Korea, the most prevalent being candidiasis, *M. tuberculosis* infection, *P. jirovecii* pneumonia, CMV infection, and herpes zoster, while toxoplasmosis and HIV-related malignancy, such as Kaposi's sarcoma, were rare. A major predictor of OD was a low CD4 T cell count at the time of HIV diagnosis. Based on our data, strategies to diagnose HIV at an early stage of infection, and timely antiretroviral therapy, are needed to reduce ODs among HIV-infected patients in Korea.

KEY MESSAGE

1. Candidiasis, *Mycobacterium tuberculosis* infection, *Pneumocystis jirovecii* pneumonia, cytomegalovirus infection, and herpes zoster were common opportunistic diseases among Korean human immunodeficiency virus (HIV)-infected patients.
2. Toxoplasmosis and HIV-related malignancy, such as lymphoma, Kaposi's sarcoma, and cervical carcinoma, were rare in Korea.
3. A major predictor of opportunistic diseases was a low CD4 T cell count at the time of HIV diagnosis.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

This study was supported by a fund of the Chronic Infectious Disease Cohort Study (Korea HIV/AIDS Cohort Study [grant number: 4800-4859-304-260, 2013-E51006-02]) from the Korea Centers for Disease Control and Prevention. We thank the members of the HIV Korean cohort. The Korea HIV/AIDS cohort study group consists of Moon Won Kang, Min Ja Kim, Jun Hee Woo, Sang Il Kim, Youn Jeong Kim, Dae Won Park, Won Suk Choi, Jang Wook Sohn, Seong Han Kim, Seong-Heon Wie, Ji-An Hur, Yeon Joon Park, Shin-Woo Kim, Hyun-Ha Chang, Yoo Joo Kim, Joon Young Song, Joong Shik Eom, Jin Seo Lee, Jacob Lee, Hye Won Jeong, Jin Soo Lee, Hee Jung Choi, Seung Soon Lee, June Myung Kim, Jun Yong Choi, Sang Hoon Han, Nam Su Ku, Jin Young Ahn, Hyo-Youl Kim, Young Keun Kim, Yong Kyun Cho, Yoon Soo Park, Seung Kwan Lim, Young Hwa Choi, Choi Bo Youl, Hee Suk Park, Mee-Kyng Kee, Joo Shil Lee, and Sung Soon Kim.

REFERENCES

- World Health Organization. Global health observatory data repository [Internet]. Geneva: World Health Organization, 2015 [cited 2015 Sep 22]. Available from: <http://apps.who.int/ghodata>.
- Buchacz K, Baker RK, Palella FJ Jr, et al. AIDS-defining opportunistic illnesses in US patients, 1994-2007: a cohort study. *AIDS* 2010;24:1549-1559.
- Szwarcz L, Chen MJ, Vittinghoff E, Hsu L, Szwarcz S. Declining incidence of AIDS-defining opportunistic illnesses: results from 16 years of population-based AIDS surveillance. *AIDS* 2013;27:597-605.
- Kim JM, Cho GJ, Hong SK, et al. Epidemiology and clinical features of HIV infection/AIDS in Korea. *Yonsei Med J* 2003;44:363-370.
- Oh MD, Park SW, Kim HB, et al. Spectrum of opportunistic infections and malignancies in patients with human immunodeficiency virus infection in South Korea. *Clin Infect Dis* 1999;29:1524-1528.
- The Korean Society for AIDS. Clinical guidelines for the treatment and prevention of opportunistic infections in HIV infected Koreans. *Infect Chemother* 2012;44:93-139.
- Joint Committee for the Development of Korean Guideline for Tuberculosis. Korean guidelines for tuberculosis. 1st ed. Cheongju: Korea Centers for Disease Control and Prevention, 2011.
- Seage GR 3rd, Losina E, Goldie SJ, Paltiel AD, Kimmel AD, Freedberg KA. The relationship of preventable opportunistic infections, HIV-1 RNA, and CD4 cell counts to chronic mortality. *J Acquir Immune Defic Syndr* 2002;30:421-428.
- Egger M, Hirschel B, Francioli P, et al. Impact of new antiretroviral combination therapies in HIV infected patients in Switzerland: prospective multicentre study. Swiss HIV Cohort Study. *BMJ* 1997;315:1194-1199.
- Damtie D, Yismaw G, Woldeyohannes D, Anagaw B. Common opportunistic infections and their CD4 cell correlates among HIV-infected patients attending at antiretroviral therapy clinic of Gondar University Hospital, Northwest Ethiopia. *BMC Res Notes* 2013;6:534.
- Xiao J, Gao G, Li Y, et al. Spectrums of opportunistic infections and malignancies in HIV-infected patients in tertiary care hospital, China. *PLoS One* 2013;8:e75915.
- Hanna DB, Gupta LS, Jones LE, Thompson DM, Kellerman SE, Sackoff JE. AIDS-defining opportunistic illnesses in the HAART era in New York City. *AIDS Care* 2007;19:264-272.
- Narain JP, Lo YR. Epidemiology of HIV-TB in Asia. *Indian J Med Res* 2004;120:277-289.
- Ledergerber B, Egger M, Opravil M, et al. Clinical progression and virological failure on highly active antiretroviral therapy in HIV-1 patients: a prospective cohort study. Swiss HIV Cohort Study. *Lancet* 1999;353:863-868.
- Nagai H. Factors for the onset of and the exacerbation of tuberculosis. 5. The infection and prognosis of tuberculosis among patients with immunodeficiency, especially HIV-infected patients. *Kekkaku* 1999;74:753-758.
- Moreno S, Jarrin I, Iribarren JA, et al. Incidence and risk factors for tuberculosis in HIV-positive subjects by HAART status. *Int J Tuberc Lung Dis* 2008;12:1393-1400.
- Sterling TR, Lau B, Zhang J, et al. Risk factors for tuberculosis after highly active antiretroviral therapy initiation in the United States and Canada: implications for tuberculosis screening. *J Infect Dis* 2011;204:893-901.
- Seyler C, Toure S, Messou E, Bonard D, Gabillard D, Anglaret X. Risk factors for active tuberculosis after antiret-

- roviral treatment initiation in Abidjan. *Am J Respir Crit Care Med* 2005;172:123-127.
19. de Sanjose S, Mbisa G, Perez-Alvarez S, et al. Geographic variation in the prevalence of Kaposi sarcoma-associated herpesvirus and risk factors for transmission. *J Infect Dis* 2009;199:1449-1456.
 20. Manosuthi W, Chaovavanich A, Tansuphaswadikul S, et al. Incidence and risk factors of major opportunistic infections after initiation of antiretroviral therapy among advanced HIV-infected patients in a resource-limited setting. *J Infect* 2007;55:464-469.
 21. Cha HH, Lee SH, Lee DH, et al. Degree of disease progression in homeless HIV/AIDS patients during the first medical visit. *Infect Chemother* 2011;43:198-202.
 22. Chattopadhyay A, Patton LL. Smoking as a risk factor for oral candidiasis in HIV-infected adults. *J Oral Pathol Med* 2013;42:302-308.
 23. Chattopadhyay A, Patton LL. Risk indicators for HIV-associated jointly occurring oral candidiasis and oral hairy leukoplakia. *AIDS Patient Care STDS* 2007;21:825-832.