

Intussusception caused by small bowel metastasis from hepatocellular carcinoma

Young Mi Hong¹, Jeong Hee Lee², and Ki Tae Yoon¹

¹Department of Internal Medicine, Pusan National University Yangsan Hospital, Pusan National University College of Medicine, Yangsan;

²Department of Pathology, Pusan National University Yangsan Hospital, Pusan National University College of Medicine, Yangsan, Korea




Figure 1. Computed tomography (CT) imaging. (A, B) Contrast-enhanced CT showed polypoid enhancing masses (white arrow) in the distal ileum.

Figure 2. Computed tomography (CT) imaging. (A-C) Contrast-enhanced CT showed polypoid enhancing masses in the distal ileum at intussusciptions (white arrow) and dilatation of upstream small bowel dilatation (blue arrow).

Figure 3. Histologic findings. (A) Left lower part of the figure showed normal mucosa of the small intestine (H&E, $\times 20$) and (B) upper part showed metastatic hepatocellular carcinoma which stains positively for hepatocyte specific antigen (DAKO 1:50, $\times 20$). (C) The tumor cells are arranged in trabecular and tubular patterns with mild nuclear atypia (H&E stain, $\times 200$). (D) The immunohistochemistry of glypican-3 (cell marquer, 1:200) shows positive reaction for tumor cells of the same field, not for intestinal mucosal epithelia (right lower of the figure, $\times 200$).

A 75-year-old man with chronic hepatitis B was diagnosed with pulmonary metastases from hepatocellular carcinoma (HCC) and was treated in order with sorafenib, regorafenib and nivolumab. However, nivolumab treatment was discontinued after the 13th administration because of poor general condition. After 2 years of conservative management, follow-up computed tomography (CT) showed no definite viable tumor in the liver and no pulmonary metastatic lesions and tumor markers were not elevated. However, iron deficiency anemia (IDA) was observed. Fecal occult blood test was negative. Although further esophagogastroduodenoscopy and colonoscopy were performed, no definite cause of the gastrointestinal (GI) bleeding was identified. While prescribing iron supplements, we performed follow-up examinations. Six months later, follow-up abdominal CT showed polypoid enhancing masses in the distal ileum (Fig. 1). There was no viable intrahepatic lesion. Laboratory results showed IDA and elevated protein induced by vitamin K absence or antagonist-II level (90.8 mAU/mL). The patient denied melena. Polypoid lesions were not detected on colonoscopy and video capsule endoscopy. Three months later, the patient visited the emergency room complaining of nausea, vomiting and abdominal pain. The AFP level was 1.9 ng/mL and PIVKA-II level was 98.7 mAU/mL. An abdominal CT scan showed polypoid enhancing masses in the distal ileum at intussusciens, suggesting intussusception of the ileum with a lead point (Fig. 2). Segmental small bowel resection and end-to end anastomosis were performed. The histological findings were consistent with those of metastatic HCC (Fig. 3). Small bowel intussusception in adult due to metastasis from HCC is extremely rare. To date, only four cases have been reported. Two cases were diagnosed with HCC metastasis to the small bowel manifesting as intussusception, and one

case was diagnosed with double-balloon enteroscopy, and the other case was diagnosed with ^{11}C -Acetate PET/CT. Our case was a small bowel metastasis of HCC manifested as obscure-occult GI bleeding, resulting in intussusception. To our knowledge, this is the first report of HCC metastasis to the small bowel manifested as obscure-occult GI bleeding. Even if HCC metastasis to the small bowel is rare, if HCC patients exhibit obscure-occult GI bleeding symptoms, it should be considered.

This study was approved by the Institutional Review Board of Pusan National University Yangsan Hospital (IRB No. 05-2023-039). Requirement for informed consent was waived after review of IRB because it was practically impossible and this study was of retrospective design.

Received : March 24, 2023

Revised : May 10, 2023

Accepted : May 18, 2023

Correspondence to

Ki Tae Yoon, M.D., Ph.D.

Department of Internal Medicine, Pusan National University Yangsan Hospital, Pusan National University College of Medicine, 20 Geumo-ro, Mulgeum-eup, Yangsan 50612, Korea

Tel: +82-55-360-2362, Fax: +82-55-360-2154

E-mail: ktyoon@pusan.ac.kr

<https://orcid.org/0000-0002-8580-0239>

CRedit authorship contributions

Young Mi Hong: conceptualization, writing - original draft, writing - review & editing; Jeong Hee Lee: resources, data curation; Ki Tae Yoon: writing - review & editing

Conflicts of interest

The authors disclose no conflicts.

Funding

This study was supported by 2023 research grant from Pusan National University Yangsan Hospital.