

Supplementary Table 1. Annual numbers and percentages of the study patients classified by sex, use of insulin, age, treatment department, and duration of medication

Variable	Year					
	2009	2010	2011	2012	2013	2014
Sex						
Male	3,806 (50.16)	4,030 (49.95)	4,422 (50.42)	4,405 (50.34)	4,433 (51.11)	4,216 (51.95)
Female	3,782 (49.84)	4,038 (50.05)	4,348 (49.58)	4,346 (49.66)	4,241 (48.89)	3,900 (48.05)
Use of insulin						
Yes	1,839 (24.24)	1,929 (23.91)	2,037 (23.23)	2,026 (23.15)	1,902 (21.93)	1,640 (20.21)
No	5,749 (75.76)	6,139 (76.09)	6,733 (76.77)	6,725 (76.85)	6,772 (78.07)	6,476 (79.79)
Age, yr						
< 40	285 (3.76)	304 (3.77)	304 (3.47)	283 (3.23)	271 (3.12)	241 (2.97)
40 to < 50	745 (9.82)	744 (9.22)	747 (8.52)	706 (8.07)	681 (7.85)	634 (7.81)
50 to < 60	1,863 (24.55)	1,970 (24.42)	2,167 (24.71)	2,126 (24.29)	1,970 (22.71)	1,826 (22.50)
60 to < 70	2,761 (36.39)	2,834 (35.13)	3,074 (35.05)	2,929 (33.47)	2,837 (32.71)	2,591 (31.92)
70 to < 80	1,637 (21.57)	1,883 (23.34)	2,087 (23.80)	2,284 (26.10)	2,453 (28.28)	2,358 (29.05)
≥ 80	297 (3.91)	333 (4.13)	391 (4.46)	423 (4.83)	462 (5.33)	466 (5.74)
Treatment department						
Endo	1,727 (22.76)	1,901 (23.56)	2,004 (22.85)	2,027 (23.16)	2,019 (23.28)	2,025 (24.95)
Endo & others	2,841 (37.44)	2,996 (37.13)	3,274 (37.33)	3,217 (36.76)	3,112 (35.88)	2,708 (33.37)
Others	3,020 (39.80)	3,171 (39.30)	3,492 (39.82)	3,507 (40.08)	3,543 (40.85)	3,383 (41.68)
Duration of medication, yr						
< 3	2,315 (30.51)	2,356 (29.20)	2,497 (28.47)	2,216 (25.32)	1,957 (22.56)	1,811 (22.31)
3–5	2,173 (28.64)	2,323 (28.79)	2,405 (27.42)	2,386 (27.27)	2,410 (22.78)	2,158 (26.59)
≥ 6	3,100 (40.85)	3,389 (42.01)	3,868 (42.01)	4,149 (47.41)	4,307 (49.65)	4,147 (51.10)
Total	7,588 (100)	8,068 (100)	8,770 (100)	8,751 (100)	8,674 (100)	8,116 (100)

Values are presented as number (%).