

Supplementary Table 6. Association between food intake and the risk of symptom at postlaparoscopic cholecystectomy by multivariable logistic regression analysis

Variable	Quartile of dietary intake			p for trend ^a
	Q1	Q2	Q3	
Grain, g				
Number of S/A	13/8	3/8	19/8	
Cut-off	≤ 253.7	253.7 < to ≤ 346.5	> 346.5	0.873
OR (95% CI) ^b	1	0.185 (0.03–1.05)	1.050 (0.23–4.76)	
Potato, g				
Number of S/A	7/8	17/8	11/8	
Cut-off	≤ 13.2	13.2 < to ≤ 47.1	> 47.1	0.649
OR (95% CI)	1	3.008 (0.71–12.75)	1.312 (0.29–5.94)	
Legume, g				
Number of S/A	12/8	14/8	9/8	
Cut-off	≤ 38.4	38.4 < to ≤ 122.8	> 122.8	0.461
OR (95% CI)	1	0.732 (0.18–2.91)	0.550 (0.13–2.40)	
Vegetable, g				
Number of S/A	6/8	11/8	18/8	
Cut-off	≤ 126.3	126.3 < to ≤ 222.3	> 222.3	0.434
OR (95% CI)	1	2.034 (0.45–9.16)	2.223 (0.46–10.83)	
Fruit, g				
Number of S/A	8/8	7/8	20/8	
Cut-off	≤ 120.4	120.4 < to ≤ 187.2	> 187.2	0.229
OR (95% CI)	1	0.667 (0.14–3.09)	1.690 (0.43–6.70)	
Meat, g				
Number of S/A	17/8	8/8	10/8	
Cut-off	≤ 50.7	50.7 < to ≤ 129.3	> 129.3	0.274
OR (95% CI)	1	0.369 (0.09–1.54)	0.361 (0.08–1.69)	
Egg, g				
Number of S/A	8/9	11/7	16/8	
Cut-off	≤ 15.2	15.2 < to ≤ 31.2	> 31.2	0.382
OR (95% CI)	1	1.159 (0.24–5.49)	1.788 (0.43–7.36)	
Fish and seafood, g				
Number of S/A	14/8	13/8	8/8	
Cut-off	≤ 40.6	40.6 < to ≤ 88.6	> 88.6	0.355
OR (95% CI)	1	1.205 (0.32–4.60)	0.503 (0.10–2.42)	
Dairy product, g				
Number of S/A	12/8	10/8	13/8	
Cut-off	≤ 56.9	56.9 < to ≤ 174.6	> 174.6	0.609
OR (95% CI)	1	0.413 (0.09–1.98)	1.195 (0.29–4.94)	
Beverage, g				
Number of S/A	16/8	11/8	8/8	
Cut-off	≤ 177.1	177.1 < to ≤ 423.6	> 423.6	0.311
OR (95% CI)	1	0.822 (0.20–3.31)	0.428 (0.09–2.15)	
Others, g				
Number of S/A	17/8	9/8	9/8	
Cut-off	≤ 20.9	20.9 < to ≤ 57.1	> 57.1	0.249
OR (95% CI)	1	0.417 (0.10–1.71)	0.332 (0.07–1.48)	

S/A, symptomatic/asymptomatic; OR, odds ratio; CI, confidence interval.

^aEstimates of p values for a linear trend were based on linear scores derived from the medians of quartiles for intake of nutrients among asymptomatic patients.

^bOR was adjusted for total energy intake and medical of digestive system disease.