

Supplementary Table 1. Demographic and clinical characteristics and outcome of patients with scrub typhus myocarditis

Age, yr	Sex	ER		F/U		CAD	CRP	NT pro-BNP, pg/mL	Peak CK, IU/L	Peak CK-MB, IU/L	Peak TnT, ng/mL	Initial echocardiogram	Initial PE	F/U LVEF	Cause of death	
		ECG STE	TWI	ECG STE	TWI											
70	M	AF	1	0	SR	0	0	22.6	29	3.5	0.4	Hypokinesia at anterior, anteroseptal wall	44	1	NA	-
87	F	AF	1	0	SR	0	0	1.0	543	7.9	0.55	Diffuse global hypokinesia	14	0	Improved 62%	-
76	F	AF	0	0	SR	0	0	5.7	34	2	0.02	Diffuse global hypokinesia	23	1	NA	-
84	F	AF	1	0	SR	0	0	3.7	698	28.3	0.22	Diffuse LV wall swelling	69	0	NA	-
68	M	AF	1	0	SR	0	0	20.7	197	2.3	0.01	Hypokinesia at anterolateral, posterolateral wall	28	0	Improved 43%	-
55	F	SR	0	1	SR	0	1	0.1	112	3.8	0.03	Diffuse global hypokinesia	19	1	Improved 71%	-
61	M	SR	1	0	SR	0	0	21.7	907	17.6	1.8	Hypokinesia at anterolateral, posterolateral wall	34	1	Expired	VF
51	F	SR	1	1	SR	0	1	0.8	315,000	7.1	0.5	Diffuse global hypokinesia	26	1	Expired	VF
46	M	SR	1	1	SR	0	1	9.3	194	2.7	0.05	Diffuse global hypokinesia	27	0	Improved 72%	-
67	M	SR	1	0	SR	0	0	0.1	2,511	10.4	0.02	Diffuse global hypokinesia	22	0	NA	-
83	F	SR	1	0	SR	0	0	24.3	449	19.1	0.5	Apical ballooning	35	0	NA	-
81	M	SR	1	0	SR	0	0	25.5	182	4.4	0.03	Diffuse global hypokinesia	24	1	NA	-
79	F	SR	1	0	SR	0	0	3.7	NA	2.9	0.05	Diffuse global hypokinesia	43	0	NA	-

ER, emergency room; ECG, electrocardiography; STE, ST-segment elevation; TWI, T-wave inversion; F/U, follow-up; CAD, coronary artery disease; CRP, C-reactive protein; NT pro-BNP, N-terminal pro-hormone of brain natriuretic peptide; CK, creatinine kinase; CK-MB, creatinine kinase-MB; TnT, troponin T; LVEF, left ventricular ejection fraction; PE, pericardial effusion; AF, atrial fibrillation; SR, sinus rhythm; NA, not applicable; LV, left ventricular; VF, ventricular fibrillation.