

Supplementary Table 3. Baseline laboratory findings of patients with scrub typhus with and without atrial fibrillation admitted to the intensive care unit

Variable	All patients	Without AF (n = 68)	With AF (n = 21)	p value
C-reactive protein, mg/dL	11.1 ± 7.9	11.4 ± 7.9	12.1 ± 8.0	0.813
White blood cells, × 10 ³ /μL	10.8 ± 9.6	7.9 ± 10.1	12.1 ± 4.8	0.328
AST, IU/L	102.6 ± 126.8	100.1 ± 138.8	107.7 ± 93.0	0.813
ALT, IU/L	81.2 ± 125.7	81.7 ± 140.0	65.5 ± 56.1	0.606
Total bilirubin, mg/dL	1.0 ± 1.7	1.3 ± 2.0	0.9 ± 0.6	0.390
GGTP, IU/L	113.9 ± 169.0	123.8 ± 185.8	87.4 ± 117.7	0.401
ALP, IU/L	138.7 ± 110.2	137.9 ± 107.4	135.6 ± 108.3	0.930
Albumin	3.2 ± 0.6	3.2 ± 0.6	2.8 ± 0.5	0.011 ^a
Protein	5.2 ± 20.8	6.4 ± 0.8	6.0 ± 0.8	0.049 ^a
LDH, IU/L	898.1 ± 875.0	940.9 ± 1,016.3	817.8 ± 316.9	0.605
BUN, mg/dL	27.6 ± 15.7	27.7 ± 15.0	32.6 ± 17.6	0.211
Creatinine, mg/dL	1.3 ± 1.2	1.4 ± 1.4	1.2 ± 0.5	0.603
Prothrombin time, sec	14.7 ± 1.4	14.6 ± 1.3	15.2 ± 1.9	0.115

Values are presented as mean ± SD.

AF, atrial fibrillation; AST, aspartate aminotransferase; ALT, alanine transaminase; GGTP, γ-glutamyl transpeptidase; ALP, alkaline phosphatase; LDH, lactate dehydrogenase; BUN, blood urea nitrogen.

^aStatistically significant.