

Supplementary Table 2. Body fluid distribution according to the diagnosis of AHF among non-dyspnea, AHF, and non-AHF group

Variable	Non-dyspnea (n = 50)	AHF (n = 50)	Non-AHF (n = 50)
Weight, kg	63.1 ± 12.1	62.6 ± 13.0	63.0 ± 11.0
TBW, L	33.3 ± 7.5	33.7 ± 8.3	32.0 ± 6.6
ICW, L	20.4 ± 4.9	19.8 ± 5.0	19.6 ± 4.6
ECW, L	12.9 ± 2.7	13.9 ± 3.5	12.3 ± 2.3
ECW/TBW	0.389 ± 0.013	0.412 ± 0.017 ^a	0.388 ± 0.023
ECW/TBW of upper extremities	0.382 ± 0.007	0.388 ± 0.013 ^b	0.380 ± 0.007
ECW/TBW of trunk	0.389 ± 0.014	0.413 ± 0.017 ^a	0.389 ± 0.020
ECW/TBW of lower extremities	0.392 ± 0.016	0.419 ± 0.020 ^a	0.390 ± 0.026

Values are presented as mean ± standard deviation.

AHF, acute heart failure; TBW, total body water; ICW, intracellular water; ECW, extracellular water.

^a $p < 0.001$ compared with non-dyspnea group.

^b $p < 0.05$ compared with non-dyspnea group.