

Supplementary Table 1. Estimated clinical primary patency and target lesion revascularization rate with 95% CI

Variable	Non-CTO-POBA	Non-CTO-Stenting	CTO-POBA	CTO-Stenting
Estimated clinical primary patency rate with 95% CI (%)				
1 yr	84.2 (80.1–88.5)	89.6 (86.6–92.8)	83.7 (79.2–88.5)	83.9 (80.5–87.5)
2 yr	79.5 (74.7–84.6)	84.5 (80.7–88.6)	76.1 (70.5–82.1)	77.1 (72.9–81.5)
3 yr	74.7 (68.4–81.5)	80.4 (75.6–85.5)	66.8 (58.7–76.1)	67.4 (61.6–73.7)
4 yr	69.3 (61.4–78.3)	77.7 (72.2–83.6)	64.2 (55.2–74.7)	62.7 (55.9–70.4)
5 yr	65.0 (55.8–75.6)	71.2 (64.1–79.2)	64.2 (55.2–74.7)	61.3 (54.2–69.3)
Estimated target lesion revascularization rate with 95% CI (%)				
1 yr	88.5 (84.8–92.3)	93.0 (90.4–95.7)	89.9 (86.2–93.9)	89.0 (86.0–92.0)
2 yr	82.8 (78.2–87.7)	89.7 (86.4–93.1)	85.8 (81.2–90.7)	82.5 (78.7–86.5)
3 yr	80.4 (74.9–86.3)	86.1 (81.7–90.7)	76.9 (69.2–85.3)	73.6 (68.0–79.6)
4 yr	78.7 (72.5–85.4)	82.4 (77.0–88.1)	74.8 (66.6–84.1)	68.1 (61.4–75.6)
5 yr	74.4 (66.5–83.3)	74.9 (67.6–82.9)	74.8 (66.6–84.1)	65.4 (58.1–73.6)

CI, confidence interval; CTO, chronic total occlusion; POBA, plain old balloon angioplasty.