

Supplementary Table 2. Patients' baseline characteristics in the whole population

Characteristic	POBA (n = 528)	Stenting (n = 801)	p value
Age	68.68 ± 8.68	69.73 ± 9.73	0.043
Male sex	392 (74.2)	656 (81.9)	0.001
BMI	23.80 ± 3.80	23.28 ± 3.28	0.017
Hypertension	414 (78.4)	606 (75.7)	0.273
DM	364 (68.9)	464 (57.9)	< 0.001
Hypercholesterolemia	209 (39.6)	330 (41.2)	0.596
Current smoker	121 (22.9)	268 (33.5)	< 0.001
CHF	40 (7.6)	57 (7.1)	0.836
CAD	313 (59.3)	454 (56.7)	0.377
CKD	157 (29.7)	150 (18.7)	< 0.001
Previous stroke	94 (17.8)	117 (14.6)	0.138
Previous bypass surgery	56 (10.6)	61 (7.6)	0.074
Previous amputation	60 (11.4)	32 (4.0)	< 0.001
Previous PTA	83 (15.7)	97 (12.1)	0.072
CLI	233 (44.1)	238 (29.7)	< 0.001
Aspirin	449 (87.2)	687 (87.3)	> 0.999
Clopidogrel	413 (80.2)	689 (87.5)	< 0.001
Cilostazol	212 (41.2)	335 (42.6)	0.657
Statin	361 (70.1)	574 (72.9)	0.294

Values are presented as mean ± SD or number (%).

POBA, plain old balloon angioplasty; BMI, body mass index; DM, diabetes mellitus; CHF, congestive heart failure; CAD, coronary artery disease; CKD, chronic kidney disease; PTA, percutaneous transluminal angioplasty; CLI, critical limb ischemia.