

Supplementary Table 6. Angina symptoms grade and calcium channel blocker by ordered logistic regression in patients with definite spasm

Variable	Unadjusted			Adjusted		
	OR	95% CI	p value	OR	95% CI	p value
1st year						
Diltiazem	Reference			Reference		
Amlodipine	1.33	0.71–2.49	0.367	1.13	0.59–2.16	0.718
Nifedipine	0.49	0.27–0.90	0.021	0.38	0.20–0.72	0.003
Benidipine	0.64	0.25–1.62	0.349	0.53	0.20–1.38	0.194
2nd year						
Diltiazem	Reference			Reference		
Amlodipine	1.46	0.65–3.29	0.364	1.17	0.50–2.75	0.711
Nifedipine	0.34	0.18–0.65	0.001	0.29	0.15–0.56	0.000
Benidipine	0.28	0.11–0.74	0.010	0.20	0.07–0.55	0.002
3rd year						
Diltiazem	Reference			Reference		
Amlodipine	0.85	0.35–2.04	0.712	0.78	0.31–1.94	0.589
Nifedipine	0.35	0.18–0.66	0.001	0.27	0.14–0.54	0.000
Benidipine	0.15	0.06–0.39	0.000	0.13	0.05–0.36	0.000

Adjusted models included age, sex, smoking, hypertension, diabetes mellitus, spasm severity, degree of atherosclerosis by angiography, history of coronary heart disease.

OR, odds ratio; CI, confidence interval.