

Supplementary Table 5. Sensitivity analyses^a

Sensitivity analyses	OR (95% CI)
Excluding participants without fasting samples (n = 398)	
Per 1 SD higher of relative change in WC	1.41 (1.16–1.72)
Per 1 SD higher of relative change in WHtR	1.38 (1.13–1.67)
Per 1 SD higher of relative change in BRI	1.36 (1.13–1.64)
Excluding participants using drugs for hypertension (n = 379)	
Per 1 SD higher of relative change in WC	1.36 (1.12–1.66)
Per 1 SD higher of relative change in WHtR	1.35 (1.11–1.65)
Per 1 SD higher of relative change in BRI	1.36 (1.12–1.66)
Excluding participants using drugs for dyslipidemia (n = 85)	
Per 1 SD higher of relative change in WC	1.42 (1.19–1.69)
Per 1 SD higher of relative change in WHtR	1.40 (1.18–1.70)
Per 1 SD higher of relative change in BRI	1.40 (1.18–1.65)

OR, odds ratio; CI, confidence interval; SD, standard deviation; WC, waist circumference; WHtR, waist-height ratio; BRI, body roundness index.

^aAdjusted for age, sex, body mass index, history of smoking and drinking, systolic blood pressure, hemoglobin A1c, total cholesterol/high-density lipoprotein cholesterol, uric acid, C-reactive protein, and walking speed at baseline.