

Supplementary Table 1. Comparison of survey results according to the respondents' affiliations

Variable	Affiliation			p value
	Primary care clinic	Community hospital	University hospital	
Need for antibiotic treatment in uncomplicated diverticulitis (n = 193)				0.260
Yes	18/21 (85.7)	52/55 (94.5)	101/117 (86.3)	
No	3/21 (14.3)	3/55 (5.5)	16/117 (13.7)	
Suggested duration of antibiotic treatment, d (n = 185)				0.500
≤ 1	0/20 (0.0)	0/52 (0.0)	1/113 (0.9)	
2–3	0/20 (0.0)	5/52 (9.6)	5/113 (4.4)	
4–7	12/20 (60.0)	35/52 (67.3)	77/113 (68.1)	
> 7	8/20 (40.0)	12/52 (23.1)	30/113 (26.5)	
Need for bowel rest in uncomplicated diverticulitis (n = 190)				0.078
Yes	7/21 (33.3)	33/53 (62.3)	61/116 (52.6)	
No	14/21 (66.7)	20/53 (37.7)	55/116 (47.4)	
Need for emergency surgery in diverticulitis with an abscess or microperforation (n = 189)				< 0.001
Yes	6/20 (30.0)	4/53 (7.5)	4/116 (3.4)	
No	14/20 (70.0)	49/53 (92.5)	112/116 (96.6)	
Treatment for recurrent disease (n = 185)				0.277
Favor repeated medical treatment	6/20 (30.0)	13/52 (25.0)	23/113 (20.4)	
Favor surgical treatment	0/20 (0.0)	1/52 (1.9)	10/113 (8.8)	
Favor individualized	14/20 (70.0)	38/52 (73.1)	80/113 (70.8)	
Elective surgery recommendation for third recurrence (n = 185)				0.056
Yes	8/20 (40.0)	11/52 (21.2)	45/113 (39.8)	
No	12/20 (60.0)	41/52 (78.8)	68/113 (60.2)	
Elective surgery recommendation for fourth recurrence (n = 185)				0.031
Yes	9/20 (45.0)	28/52 (53.8)	79/113 (69.9)	
No	11/20 (55.0)	24/52 (46.2)	34/113 (30.1)	

Values are presented with number of responses per total number of respondents (%).