

Supplementary Table 1. Laboratory findings at the time of RA and suspected NTM-PD

Variable	At RA diagnosis	At suspected NTM	<i>p</i> value
ESR, mm/h	46.0 ± 30.2	37.6 ± 30.6	0.014
CRP, mg/dL	2.37 ± 3.46	3.04 ± 5.64	0.329
DAS28	5.0 ± 1.6	3.7 ± 1.3	< 0.001
WBC, / μ L	8,090.6 ± 2,860.9	7,447.6 ± 2,894.1	0.110
Neutrophil, / μ L	5,524.8 ± 2,537.9	5,199.7 ± 3,082.6	0.420
Lymphocyte, / μ L	1,905.3 ± 745.5	1,547.8 ± 674.9	< 0.001
Hb, mg/dL	13.5 ± 12.3	12.2 ± 1.8	0.335
Hct, %	36.7 ± 4.0	36.7 ± 5.2	0.936
Platelet, $\times 10^3$ / μ L	298.3 ± 108.9	266.5 ± 95.0	0.004
Total protein, g/dL	7.2 ± 0.8	7.0 ± 0.8	0.003
Albumin, g/dL	4.1 ± 0.4	4.1 ± 0.6	0.361

CRP, C-reactive protein; DAS, disease activity score; ESR, erythrocyte sedimentation rate; Hb, hemoglobin; Hct, hematocrit; NTM-PD, non-tuberculous mycobacterial pulmonary disease; RA, rheumatoid arthritis; WBC, white blood cell.
p values are calculated using chi-square test or Student's t-test.