

Supplementary Table 3. The results of node-splitting method for local inconsistencies

Comparison	Direct	Indirect	Difference	<i>p</i> value
48VR				
BSV vs. ETV	-0.104 (-1.040 to 0.832)	-0.146 (-0.644 to 0.353)	0.042 (-1.019 to 1.103)	0.938
BSV vs. TAF	0.220 (-0.325 to 0.765)	0.318 (-0.372 to 1.009)	-0.098 (-0.978 to 0.781)	0.827
BSV vs. TDF	0.224 (-0.449 to 0.896)	0.154 (-0.400 to 0.709)	0.069 (-0.862 to 0.941)	0.876
ETV vs. TDF	0.319 (0.172 to 0.467)	0.277 (-0.773 to 1.328)	0.042 (-1.019 to 1.103)	0.938
TAF vs. TDF	-0.097 (-0.515 to 0.320)	0.001 (-0.773 to 0.775)	-0.098 (-0.970 to 0.773)	0.827
96VR				
BSV vs. TAF	-0.405 (-2.045 to 1.236)	-0.403 (-1.902 to 1.096)	-0.003 (-2.226 to 2.220)	0.998
ETV vs. TDF	-0.490 (-1.977 to 0.997)	-0.492 (-2.145 to 1.160)	0.002 (-2.220 to 2.225)	0.998
TAF vs. TDF	-0.087 (-0.280 to 0.106)	-0.084 (-0.278 to 0.109)	-0.003 (-2.226 to 2.221)	0.998
48BR				
BSV vs. ETV	0.965 (-0.410 to 2.340)	0.366 (-0.306 to 1.037)	0.600 (-0.931 to 2.130)	0.443
BSV vs. TAF	0.319 (-0.404 to 1.041)	0.335 (-0.643 to 1.313)	-0.016 (-1.232 to 1.200)	0.979
BSV vs. TDF	0.041 (-0.804 to 0.885)	0.364 (-0.432 to 1.161)	-0.324 (-1.484 to 0.837)	0.585
ETV vs. TDF	-0.254 (-0.505 to -0.002)	-0.853 (-2.367 to 0.660)	0.599 (-0.931 to 2.130)	0.443
TAF vs. TDF	-0.116 (-0.767 to 0.535)	-0.100 (-1.126 to 0.927)	-0.016 (-1.232 to 1.200)	0.979
48SR				
BSV vs. ETV	-0.405 (-2.047 to 1.236)	-0.403 (-1.902 to 1.096)	-0.003 (-2.226 to 2.220)	0.998
BSV vs. TDF	-0.490 (-1.977 to 0.997)	-0.492 (-2.145 to 1.160)	0.002 (-2.220 to 2.225)	0.998
ETV vs. TDF	-0.082 (-0.275 to 0.111)	-0.084 (-2.293 to 2.125)	-0.003 (-2.226 to 2.221)	0.998

Values are presented as coefficient (95% confidence interval).

48VR, 48-week virologic response; 96VR, 96-week virologic response; 48BR, 48-week biochemical response; 48SR, 48-week serologic response; BSV, besifovir dipivoxil maleate; ETV, entecavir; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate.