

Supplementary Table 2. Electrocardiogram parameters

Variable	All patients (n = 374)	Normotensive patients (n = 343)		<i>p</i> value ^{a)}	High risk patients (n = 31)	<i>p</i> value ^{b)}
		hsTnT < 60 ng/ L (n = 237)	hsTnT ≥ 60 ng/L (n = 106)			
Normal electrocardiogram	158 (42.3)	120 (50.6)	29 (27.4)	< 0.001	9 (29.0)	< 0.001
Sinus tachycardia	80 (21.4)	43 (18.1)	30 (28.3)	0.048	7 (22.6)	0.104
Atrial fibrillation	24 (6.4)	11 (4.6)	9 (8.5)	0.247	4 (12.9)	0.124
T-wave inversions in lead V1–V3	91 (24.3)	51 (21.5)	31 (29.3)	0.158	9 (29.0)	0.249
Right bundle branch block (complete or incomplete)	52 (13.9)	23 (9.7)	19 (17.9)	0.049	10 (32.3)	0.001
Complete atrioventricular block	1 (0.3)	0 (0.0)	1 (0.9)	0.309	0 (0.0)	0.282

Values are presented as number (%).

hsTnT, high-sensitivity troponin T.

^{a)}*p* value was determined through chi-squared or Mann–Whitney U test, comparing groups with hsTnT < 60 ng/L and ≥ 60 ng/L in normotensive patients.

^{b)}*p* value resulted from the ANOVA test comparing three groups.