Supplementary Table 1. Effectiveness of BDP/F compared to BUD/F in asthma

Overall (n = 24)	Changes after BDP/F	Changes after BUD/F	*p* value
Delta FEF_{25-75} (L)	-0.01 ± 0.46	-0.10 ± 0.48	0.496
Delta FEF_{25-75} (% of predicted value)	5.79 ± 38.34	-1.36 ± 14.93	0.399
Delta FEV_1 (L)	0.01 ± 0.26	0.00 ± 0.17	0.917
Delta FEV_1 (% of predicted value)	1.38 ± 12.68	0.27 ± 6.32	0.704
Delta FVC (L)	0.01 ± 0.27	0.05 ± 0.18	0.516
Delta FVC (% of predicted value)	0.58 ± 7.90	1.41 ± 4.99	0.663
Delta FEV_1/FVC (%)	0.46 ± 3.90	-0.69 ± 3.34	0.280
Delta ACQ	-0.02 ± 0.73	0.10 ± 0.78	0.578
Delta ACT	0.79 ± 3.49	-0.71 ± 3.84	0.163
Delta QLQAKA	0.05 ± 0.53	-0.17 ± 0.54	0.164

Values are presented as mean ± standard deviation using the Linear Mixed Model.
BDP/F, beclomethasone/formoterol; BUD/F, budesonide/formoterol; FEF_{25-75}, forced expiratory flow between 25% and 75% of vital capacity; FEV_1, forced expiratory volume in 1 second; FVC, forced vital capacity; ACQ: Asthma Control Questionnaire; ACT: Asthma Control Test; QLQAKA: Quality of Life Questionnaire for Adult Korean Asthmatics.