

Supplementary Table 2. OR and 95% CI for global satisfaction scores ≥ 80 based on multivariable logistic regression analyses in the patients with remission or low disease activity of rheumatoid arthritis or with no or mild disease activity of systemic lupus erythematosus

	Adjusted OR (95% CI)	
	RA (n = 101)	SLE (n = 92)
Sex		
Male	Reference	Reference
Female	0.32 (0.09–1.13)	1.16 (0.09–13.84)
Age at the survey (yr)	1.02 (0.94–1.10)	1.02 (0.95–1.09)
Duration of education (yr)		
≤ 9	Reference	Reference
9–12	3.36 (0.65–17.1)	3.84 (0.37–39.38)
≥ 12	1.87 (0.32–10.94)	8.48 (0.60–119.44)
Household monthly income		
< 300 (*10,000 won)	Reference	Reference
300–500 (*10,000 won)	9.16 (1.39–60.40)	0.53 (0.11–2.53)
≥ 500 (*10,000 won)	9.33 (1.49–58.19)	1.11 (0.14–8.35)
Current use of NSAIDs		
Yes	Reference	Reference
No	0.31 (0.05–1.65)	8.65 (1.02–73.22)
Current use of steroids		
Yes	Reference	Reference
No	1.26 (0.37–4.22)	2.39 (0.48–11.80)
Number of comorbidities	1.11 (0.57–2.15)	1.25 (0.57–2.74)

OR, odds ratio; CI, confidence interval; NSAIDs, non-steroidal anti-inflammatory drugs.